

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/128873>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

Measurements of Branching Fractions and CP -Violating Asymmetries in $B^0 \rightarrow \rho^\pm h^\mp$ Decays

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ P. Robbe,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ J. F. Kral,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ M. E. Levi,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ A. Romosan,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ D. J. Knowles,⁶ S. E. Morgan,⁶ R. C. Penny,⁶ A. T. Watson,⁶ N. K. Watson,⁶ T. Deppermann,⁷ K. Goetzen,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ K. Peters,⁷ H. Schmucker,⁷ M. Steinke,⁷ N. R. Barlow,⁸ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ C. Mackay,⁸ F. F. Wilson,⁸ K. Abe,⁹ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ P. Kyberd,¹⁰ A. K. McKemey,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Chao,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² S. McMahon,¹² R. K. Mommensen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ D. del Re,¹⁴ H. K. Hadavand,¹⁴ E. J. Hill,¹⁴ D. B. MacFarlane,¹⁴ H. P. Paar,¹⁴ Sh. Rahatlou,¹⁴ U. Schwanke,¹⁴ V. Sharma,¹⁴ J. W. Berryhill,¹⁵ C. Campagnari,¹⁵ B. Dahmes,¹⁵ N. Kuznetsova,¹⁵ S. L. Levy,¹⁵ O. Long,¹⁵ A. Lu,¹⁵ M. A. Mazur,¹⁵ J. D. Richman,¹⁵ W. Verkerke,¹⁵ T. W. Beck,¹⁶ J. Beringer,¹⁶ A. M. Eisner,¹⁶ C. A. Heusch,¹⁶ W. S. Lockman,¹⁶ T. Schalk,¹⁶ R. E. Schmitz,¹⁶ B. A. Schumm,¹⁶ A. Seiden,¹⁶ M. Turri,¹⁶ W. Walkowiak,¹⁶ D. C. Williams,¹⁶ M. G. Wilson,¹⁶ J. Albert,¹⁷ E. Chen,¹⁷ G. P. Dubois-Felsmann,¹⁷ A. Dvoretzki,¹⁷ D. G. Hitlin,¹⁷ I. Narsky,¹⁷ F. C. Porter,¹⁷ A. Ryd,¹⁷ A. Samuel,¹⁷ S. Yang,¹⁷ S. Jayatilake,¹⁸ G. Mancinelli,¹⁸ B. T. Meadows,¹⁸ M. D. Sokoloff,¹⁸ T. Abe,¹⁹ T. Barillari,¹⁹ F. Blanc,¹⁹ P. Bloom,¹⁹ P. J. Clark,¹⁹ W. T. Ford,¹⁹ U. Nauenberg,¹⁹ A. Olivas,¹⁹ P. Rankin,¹⁹ J. Roy,¹⁹ J. G. Smith,¹⁹ W. C. van Hoek,¹⁹ L. Zhang,¹⁹ J. L. Harton,²⁰ T. Hu,²⁰ A. Soffer,²⁰ W. H. Toki,²⁰ R. J. Wilson,²⁰ J. Zhang,²⁰ D. Altenburg,²¹ T. Brandt,²¹ J. Brose,²¹ T. Colberg,²¹ M. Dickopp,²¹ R. S. Dubitzky,²¹ A. Hauke,²¹ H. M. Lacker,²¹ E. Maly,²¹ R. Müller-Pfefferkorn,²¹ R. Nogowski,²¹ S. Otto,²¹ K. R. Schubert,²¹ R. Schwierz,²¹ B. Spaan,²¹ L. Wilden,²¹ D. Bernard,²² G. R. Bonneaud,²² F. Brochard,²² J. Cohen-Tanugi,²² Ch. Thiebaux,²² G. Vasileiadis,²² M. Verderi,²² A. Khan,²³ D. Lavin,²³ F. Muheim,²³ S. Playfer,²³ J. E. Swain,²³ J. Tinslay,²³ M. Andreotti,²⁴ D. Bettoni,²⁴ C. Bozzi,²⁴ R. Calabrese,²⁴ G. Cibinetto,²⁴ E. Luppi,²⁴ M. Negrini,²⁴ L. Piemontese,²⁴ A. Sarti,²⁴ E. Treadwell,²⁵ F. Anulli,^{1,*} R. Baldini-Ferrolli,²⁶ A. Calcaterra,²⁶ R. de Sangro,²⁶ D. Falciari,²⁶ G. Finocchiaro,²⁶ P. Patteri,²⁶ I. M. Peruzzi,^{26,*} M. Piccolo,²⁶ A. Zallo,²⁶ A. Buzzo,²⁷ R. Contri,²⁷ G. Crosetti,²⁷ M. Lo Vetere,²⁷ M. Macri,²⁷ M. R. Monge,²⁷ S. Passaggio,²⁷ F. C. Pastore,²⁷ C. Patrignani,²⁷ E. Robutti,²⁷ A. Santroni,²⁷ S. Tosi,²⁷ S. Bailey,²⁸ M. Morii,²⁸ M. L. Aspinwall,²⁹ W. Bhimji,²⁹ D. A. Bowerman,²⁹ P. D. Dauncey,²⁹ U. Egede,²⁹ I. Eschrich,²⁹ G. W. Morton,²⁹ J. A. Nash,²⁹ P. Sanders,²⁹ G. P. Taylor,²⁹ G. J. Grenier,³⁰ S.-J. Lee,³⁰ U. Mallik,³⁰ J. Cochran,³¹ H. B. Crawley,³¹ J. Lamsa,³¹ W. T. Meyer,³¹ S. Prell,³¹ E. I. Rosenberg,³¹ J. Yi,³¹ M. Davier,³² G. Grosdidier,³² A. Höcker,³² S. Laplace,³² F. Le Diberder,³² V. Lepeltier,³² A. M. Lutz,³² T. C. Petersen,³² S. Plaszczynski,³² M. H. Schune,³² L. Tantot,³² G. Wormser,³² V. Brigljević,³³ C. H. Cheng,³³ D. J. Lange,³³ D. M. Wright,³³ A. J. Bevan,³⁴ J. P. Coleman,³⁴ J. R. Fry,³⁴ E. Gabathuler,³⁴ R. Gamet,³⁴ M. Kay,³⁴ R. J. Parry,³⁴ D. J. Payne,³⁴ R. J. Sloane,³⁴ C. Touramanis,³⁴ J. J. Back,³⁵ P. F. Harrison,³⁵ H. W. Shorthouse,³⁵ P. Strother,³⁵ P. B. Vidal,³⁵ C. L. Brown,³⁶ G. Cowan,³⁶ R. L. Flack,³⁶ H. U. Flaecher,³⁶ S. George,³⁶ M. G. Green,³⁶ A. Kurup,³⁶ C. E. Marker,³⁶ T. R. McMahon,³⁶ S. Ricciardi,³⁶ F. Salvatore,³⁶ G. Vaitsas,³⁶ M. A. Winter,³⁶ D. Brown,³⁷ C. L. Davis,³⁷ J. Allison,³⁸ R. J. Barlow,³⁸ A. C. Forti,³⁸ P. A. Hart,³⁸ F. Jackson,³⁸ G. D. Lafferty,³⁸ A. J. Lyon,³⁸ J. H. Weatherall,³⁸ J. C. Williams,³⁸ A. Farbin,³⁹ A. Jawahery,³⁹ D. Kovalskyi,³⁹ C. K. Lae,³⁹ V. Lillard,³⁹ D. A. Roberts,³⁹ G. Blaylock,⁴⁰ C. Dallapiccola,⁴⁰ K. T. Flood,⁴⁰ S. S. Hertzbach,⁴⁰ R. Kofler,⁴⁰ V. B. Koptchev,⁴⁰ T. B. Moore,⁴⁰ S. Saremi,⁴⁰ H. Staengle,⁴⁰ S. Willocq,⁴⁰ R. Cowan,⁴¹ G. Sciolla,⁴¹ F. Taylor,⁴¹ R. K. Yamamoto,⁴¹ D. J. J. Mangeol,⁴² M. Milek,⁴² P. M. Patel,⁴² A. Lazzaro,⁴³ F. Palombo,⁴³ J. M. Bauer,⁴⁴ L. Cremaldi,⁴⁴ V. Eschenburg,⁴⁴ R. Godang,⁴⁴ R. Kroeger,⁴⁴ J. Reidy,⁴⁴ D. A. Sanders,⁴⁴ D. J. Summers,⁴⁴ H. W. Zhao,⁴⁴ C. Hast,⁴⁵ P. Taras,⁴⁵ H. Nicholson,⁴⁶ C. Cartaro,⁴⁷ N. Cavallo,⁴⁷ G. De Nardo,⁴⁷ F. Fabozzi,^{47,†} C. Gatto,⁴⁷ L. Lista,⁴⁷ P. Paolucci,⁴⁷ D. Piccolo,⁴⁷ C. Sciacca,⁴⁷ M. A. Baak,⁴⁸ G. Raven,⁴⁸ J. M. LoSecco,⁴⁹ T. A. Gabriel,⁵⁰ B. Brau,⁵¹ T. Pulliam,⁵¹ J. Brau,⁵² R. Frey,⁵² C. T. Potter,⁵² N. B. Sinev,⁵² D. Strom,⁵² E. Torrence,⁵² F. Colecchia,⁵³ A. Dorigo,⁵³ F. Galeazzi,⁵³ M. Margoni,⁵³ M. Morandin,⁵³ M. Posocco,⁵³ M. Rotondo,⁵³ F. Simonetto,⁵³ R. Stroili,⁵³ G. Tiozzo,⁵³ C. Voci,⁵³ M. Benayoun,⁵⁴ H. Briand,⁵⁴ J. Chauveau,⁵⁴ P. David,⁵⁴ Ch. de la Vaissière,⁵⁴

L. Del Buono,⁵⁴ O. Hamon,⁵⁴ M. J. J. John,⁵⁴ Ph. Leruste,⁵⁴ J. Ocariz,⁵⁴ M. Pivk,⁵⁴ L. Roos,⁵⁴ J. Stark,⁵⁴ S. T'Jampens,⁵⁴ P. F. Manfredi,⁵⁵ V. Re,⁵⁵ L. Gladney,⁵⁶ Q. H. Guo,⁵⁶ J. Panetta,⁵⁶ C. Angelini,⁵⁷ G. Batignani,⁵⁷ S. Bettarini,⁵⁷ M. Bondioli,⁵⁷ F. Bucci,⁵⁷ G. Calderini,⁵⁷ M. Carpinelli,⁵⁷ F. Forti,⁵⁷ M. A. Giorgi,⁵⁷ A. Lusiani,⁵⁷ G. Marchiori,⁵⁷ F. Martinez-Vidal,^{57,‡} M. Morganti,⁵⁷ N. Neri,⁵⁷ E. Paoloni,⁵⁷ M. Rama,⁵⁷ G. Rizzo,⁵⁷ F. Sandrelli,⁵⁷ J. Walsh,⁵⁷ M. Haire,⁵⁸ D. Judd,⁵⁸ K. Paick,⁵⁸ D. E. Wagoner,⁵⁸ N. Danielson,⁵⁹ P. Elmer,⁵⁹ C. Lu,⁵⁹ V. Miftakov,⁵⁹ J. Olsen,⁵⁹ A. J. S. Smith,⁵⁹ E. W. Varnes,⁵⁹ F. Bellini,⁶⁰ G. Cavoto,^{59,60} R. Faccini,^{14,60} F. Ferrarotto,⁶⁰ F. Ferroni,⁶⁰ M. Gaspero,⁶⁰ M. A. Mazzoni,⁶⁰ S. Morganti,⁶⁰ M. Pierini,⁶⁰ G. Piredda,⁶⁰ F. Safai Tehrani,⁶⁰ C. Voena,⁶⁰ S. Christ,⁶¹ G. Wagner,⁶¹ R. Waldi,⁶¹ T. Adye,⁶² N. De Groot,⁶² B. Franek,⁶² N. I. Geddes,⁶² G. P. Gopal,⁶² E. O. Olaiya,⁶² S. M. Xella,⁶² R. Aleksan,⁶³ S. Emery,⁶³ A. Gaidot,⁶³ S. F. Ganzhur,⁶³ P.-F. Giraud,⁶³ G. Hamel de Monchenault,⁶³ W. Kozanecki,⁶³ M. Langer,⁶³ G. W. London,⁶³ B. Mayer,⁶³ G. Schott,⁶³ G. Vasseur,⁶³ Ch. Yeché,⁶³ M. Zito,⁶³ M. V. Purohit,⁶⁴ A. W. Weidemann,⁶⁴ F. X. Yumiceva,⁶⁴ D. Aston,⁶⁵ R. Bartoldus,⁶⁵ N. Berger,⁶⁵ A. M. Boyarski,⁶⁵ O. L. Buchmueller,⁶⁵ M. R. Convery,⁶⁵ D. P. Coupal,⁶⁵ D. Dong,⁶⁵ J. Dorfan,⁶⁵ D. Dujmic,⁶⁵ W. Dunwoodie,⁶⁵ R. C. Field,⁶⁵ T. Glanzman,⁶⁵ S. J. Gowdy,⁶⁵ E. Grauges-Pous,⁶⁵ T. Hadig,⁶⁵ V. Halyo,⁶⁵ T. Hryn'ova,⁶⁵ W. R. Innes,⁶⁵ C. P. Jessop,⁶⁵ M. H. Kelsey,⁶⁵ P. Kim,⁶⁵ M. L. Kocian,⁶⁵ U. Langenegger,⁶⁵ D. W. G. S. Leith,⁶⁵ S. Luitz,⁶⁵ V. Luth,⁶⁵ H. L. Lynch,⁶⁵ H. Marsiske,⁶⁵ S. Menke,⁶⁵ R. Messner,⁶⁵ D. R. Muller,⁶⁵ C. P. O'Grady,⁶⁵ V. E. Ozcan,⁶⁵ A. Perazzo,⁶⁵ M. Perl,⁶⁵ S. Petrak,⁶⁵ B. N. Ratcliff,⁶⁵ S. H. Robertson,⁶⁵ A. Roodman,⁶⁵ A. A. Salnikov,⁶⁵ R. H. Schindler,⁶⁵ J. Schwiening,⁶⁵ G. Simi,⁶⁵ A. Snyder,⁶⁵ A. Soha,⁶⁵ J. Stelzer,⁶⁵ D. Su,⁶⁵ M. K. Sullivan,⁶⁵ H. A. Tanaka,⁶⁵ J. Va'vra,⁶⁵ S. R. Wagner,⁶⁵ M. Weaver,⁶⁵ A. J. R. Weinstein,⁶⁵ W. J. Wisniewski,⁶⁵ D. H. Wright,⁶⁵ C. C. Young,⁶⁵ P. R. Burchat,⁶⁶ A. J. Edwards,⁶⁶ T. I. Meyer,⁶⁶ C. Roat,⁶⁶ S. Ahmed,⁶⁷ M. S. Alam,⁶⁷ J. A. Ernst,⁶⁷ M. Saleem,⁶⁷ F. R. Wappler,⁶⁷ W. Bugg,⁶⁸ M. Krishnamurthy,⁶⁸ S. M. Spanier,⁶⁸ R. Eckmann,⁶⁹ H. Kim,⁶⁹ J. L. Ritchie,⁶⁹ R. F. Schwitters,⁶⁹ J. M. Izen,⁷⁰ I. Kitayama,⁷⁰ X. C. Lou,⁷⁰ S. Ye,⁷⁰ F. Bianchi,⁷¹ M. Bona,⁷¹ F. Gallo,⁷¹ D. Gamba,⁷¹ C. Borean,⁷² L. Bosisio,⁷² G. Della Ricca,⁷² S. Dittongo,⁷² S. Grancagnolo,⁷² L. Lanceri,⁷² P. Poropat,^{72,§} L. Vitale,⁷² G. Vuagnin,⁷² R. S. Panvini,⁷³ Sw. Banerjee,⁷⁴ C. M. Brown,⁷⁴ D. Fortin,⁷⁴ P. D. Jackson,⁷⁴ R. Kowalewski,⁷⁴ J. M. Roney,⁷⁴ H. R. Band,⁷⁵ S. Dasu,⁷⁵ M. Datta,⁷⁵ A. M. Eichenbaum,⁷⁵ H. Hu,⁷⁵ J. R. Johnson,⁷⁵ P. E. Kutter,⁷⁵ H. Li,⁷⁵ R. Liu,⁷⁵ F. Di Lodovico,⁷⁵ A. Mihalysi,⁷⁵ A. K. Mohapatra,⁷⁵ Y. Pan,⁷⁵ R. Prepost,⁷⁵ S. J. Sekula,⁷⁵ J. H. von Wimmersperg-Toeller,⁷⁵ J. Wu,⁷⁵ S. L. Wu,⁷⁵ and Z. Yu⁷⁵ H. Neal⁷⁶

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy³Institute of High Energy Physics, Beijing 100039, China⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom⁷Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany⁸University of Bristol, Bristol BS8 1TL, United Kingdom⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia¹²University of California at Irvine, Irvine, California 92697, USA¹³University of California at Los Angeles, Los Angeles, California 90024, USA¹⁴University of California at San Diego, La Jolla, California 92093, USA¹⁵University of California at Santa Barbara, Santa Barbara, California 93106, USA¹⁶University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA¹⁷California Institute of Technology, Pasadena, California 91125, USA¹⁸University of Cincinnati, Cincinnati, Ohio 45221, USA¹⁹University of Colorado, Boulder, Colorado 80309, USA²⁰Colorado State University, Fort Collins, Colorado 80523, USA²¹Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany²²Ecole Polytechnique, LLR, F-91128 Palaiseau, France²³University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom²⁴Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy²⁵Florida A&M University, Tallahassee, Florida 32307, USA²⁶Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy²⁷Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy²⁸Harvard University, Cambridge, Massachusetts 02138, USA

- ²⁹Imperial College London, London, SW7 2BW, United Kingdom
³⁰University of Iowa, Iowa City, Iowa 52242, USA
³¹Iowa State University, Ames, Iowa 50011-3160, USA
³²Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³³Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁴University of Liverpool, Liverpool L69 3BX, United Kingdom
³⁵Queen Mary, University of London, E1 4NS, United Kingdom
³⁶University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
³⁷University of Louisville, Louisville, Kentucky 40292, USA
³⁸University of Manchester, Manchester M13 9PL, United Kingdom
³⁹University of Maryland, College Park, Maryland 20742, USA
⁴⁰University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴¹Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴²McGill University, Montréal, Québec, Canada H3A 2T8
⁴³Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁴University of Mississippi, University, Mississippi 38677, USA
⁴⁵Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Québec, Canada H3C 3J7
⁴⁶Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁴⁷Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁴⁸NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁴⁹University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁰Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
⁵¹Ohio State University, Columbus, Ohio 43210, USA
⁵²University of Oregon, Eugene, Oregon 97403, USA
⁵³Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁴Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
⁵⁵Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
⁵⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁷Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁵⁸Prairie View A&M University, Prairie View, Texas 77446, USA
⁵⁹Princeton University, Princeton, New Jersey 08544, USA
⁶⁰Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶¹Universität Rostock, D-18051 Rostock, Germany
⁶²Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶³DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁴University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁵Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁶Stanford University, Stanford, California 94305-4060, USA
⁶⁷State University of New York, Albany, New York 12222, USA
⁶⁸University of Tennessee, Knoxville, Tennessee 37996, USA
⁶⁹University of Texas at Austin, Austin, Texas 78712, USA
⁷⁰University of Texas at Dallas, Richardson, Texas 75083, USA
⁷¹Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷²Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷³Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁴University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁵University of Wisconsin, Madison, Wisconsin 53706, USA
⁷⁶Yale University, New Haven, Connecticut 06511, USA

(Received 13 June 2003; published 13 November 2003)

We present measurements of branching fractions and CP -violating asymmetries in $B^0 \rightarrow \rho^\pm \pi^\mp$ and $B^0 \rightarrow \rho^- K^+$ decays. The results are obtained from a data sample of 88.9×10^6 $Y(4S) \rightarrow B\bar{B}$ decays collected with the $BABAR$ detector at the SLAC PEP-II asymmetric-energy B Factory. From a time-dependent maximum likelihood fit we measure the branching fractions $\mathcal{B}(B^0 \rightarrow \rho^\pm \pi^\mp) = [22.6 \pm 1.8 (\text{stat}) \pm 2.2 (\text{syst})] \times 10^{-6}$ and $\mathcal{B}(B^0 \rightarrow \rho^- K^+) = (7.3^{+1.3}_{-1.2} \pm 1.3) \times 10^{-6}$, and the CP -violating charge asymmetries $A_{CP}^{\rho\pi} = -0.18 \pm 0.08 \pm 0.03$ and $A_{CP}^{K\rho} = 0.28 \pm 0.17 \pm 0.08$, the direct CP violation parameter $C_{\rho\pi} = 0.36 \pm 0.18 \pm 0.04$ and the mixing-induced CP violation parameter $S_{\rho\pi} = 0.19 \pm 0.24 \pm 0.03$, and the dilution parameters $\Delta C_{\rho\pi} = 0.28^{+0.18}_{-0.19} \pm 0.04$ and $\Delta S_{\rho\pi} = 0.15 \pm 0.25 \pm 0.03$.

DOI: 10.1103/PhysRevLett.91.201802

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

One of the central issues in particle physics is whether the single complex phase of the Cabibbo-Kobayashi-Maskawa quark-mixing matrix [1] is sufficient to explain the observed pattern of CP violation. We present here a simultaneous measurement of branching fractions and CP -violating asymmetries in the decays $B^0 \rightarrow \rho^\pm \pi^\mp$ and $B^0 \rightarrow \rho^- K^+$ (and their charge conjugates). The *BABAR* and Belle experiments have performed searches for CP -violating asymmetries in B decays to $\pi^+ \pi^-$ [2,3], where the mixing-induced CP asymmetry is related to the angle $\alpha \equiv \arg[-V_{td}V_{tb}^*/V_{ud}V_{ub}^*]$ of the unitarity triangle as it is for $\rho^\pm \pi^\mp$. However, unlike $\pi^+ \pi^-$, $\rho^\pm \pi^\mp$ is not a CP eigenstate, and four flavor-charge configurations [$B^0(\bar{B}^0) \rightarrow \rho^\pm \pi^\mp$] must be considered [4]. Although this leads to a more complicated analysis, it benefits from a branching fraction that is nearly 5 times larger [5,6]. The extraction of α from $\rho^\pm \pi^\mp$ is complicated by the interference of decay amplitudes with different strong and weak phases. One strategy for overcoming this problem is to perform an isospin analysis including all $\rho\pi$ final states [7]. Another approach exploits the dynamical information of the full Dalitz plot to extract α and the strong amplitudes simultaneously [8].

Following a quasi-two-body approach [9], we restrict the analysis to the two regions of the $\pi^\mp \pi^0 h^\pm$ Dalitz plot ($h = \pi$ or K) that are dominated by either $\rho^+ h^-$ or $\rho^- h^+$. With $\Delta t \equiv t_{\rho h} - t_{\text{tag}}$ defined as the proper time interval between the decay of the reconstructed $B_{\rho h}^0$ and that of the other meson B_{tag}^0 , the time-dependent decay rates are given by

$$f_{Q_{\text{tag}}}^{\rho^\pm h^\mp}(\Delta t) = (1 \pm A_{CP}^{\rho h}) \frac{e^{-|\Delta t|/\tau}}{4\tau} \\ \times [1 + Q_{\text{tag}}(S_{\rho h} \pm \Delta S_{\rho h}) \sin(\Delta m_d \Delta t) \\ - Q_{\text{tag}}(C_{\rho h} \pm \Delta C_{\rho h}) \cos(\Delta m_d \Delta t)], \quad (1)$$

where $Q_{\text{tag}} = 1$ (-1) when the tagging meson B_{tag}^0 is a B^0 (\bar{B}^0), τ is the mean B^0 lifetime, and Δm_d is the $B^0 \bar{B}^0$ oscillation frequency. The time- and flavor-integrated charge asymmetries $A_{CP}^{\rho\pi}$ and $A_{CP}^{\rho K}$ measure direct CP violation. For the $\rho\pi$ mode, the quantities $S_{\rho\pi}$ and $C_{\rho\pi}$ parametrize mixing-induced CP violation related to the angle α , and flavor-dependent direct CP violation, respectively. The parameters $\Delta C_{\rho\pi}$ and $\Delta S_{\rho\pi}$ are insensitive to CP violation. $\Delta C_{\rho\pi}$ describes the asymmetry between the rates $\Gamma(B^0 \rightarrow \rho^+ \pi^-) + \Gamma(\bar{B}^0 \rightarrow \rho^- \pi^+)$ and $\Gamma(B^0 \rightarrow \rho^- \pi^+) + \Gamma(\bar{B}^0 \rightarrow \rho^+ \pi^-)$, while $\Delta S_{\rho\pi}$ is related to the strong phase difference between the amplitudes contributing to $B^0 \rightarrow \rho\pi$ decays. More precisely, one finds the relations $S_{\rho\pi} \pm \Delta S_{\rho\pi} = \sqrt{1 - (C_{\rho\pi} \pm \Delta C_{\rho\pi})^2} \sin(2\alpha_{\text{eff}}^\pm \pm \delta) \rho\pi$, where $2\alpha_{\text{eff}}^\pm = \arg[(q/p)(\bar{A}_{\rho\pi}^\pm/A_{\rho\pi}^\pm)]$, $\delta = \arg[A_{\rho\pi}^-/A_{\rho\pi}^+]$, $\arg[q/p]$ is the $B^0 \bar{B}^0$ mixing phase, and $A_{\rho\pi}^+ (\bar{A}_{\rho\pi}^+)$ and $A_{\rho\pi}^- (\bar{A}_{\rho\pi}^-)$ are

the transition amplitudes of the processes $B^0(\bar{B}^0) \rightarrow \rho^+ \pi^-$ and $B^0(\bar{B}^0) \rightarrow \rho^- \pi^+$, respectively. The angles α_{eff}^\pm are equal to α in the absence of contributions from penguin amplitudes. For the self-tagging ρK mode, the values of the four time-dependent parameters are $C_{\rho K} = 0$, $\Delta C_{\rho K} = -1$, $S_{\rho K} = 0$, and $\Delta S_{\rho K} = 0$.

The data used in this analysis were accumulated with the *BABAR* detector [10], at the SLAC PEP-II asymmetric-energy e^+e^- storage ring. The sample consists of $(88.9 \pm 1.0) \times 10^6 B\bar{B}$ pairs collected at the $Y(4S)$ resonance (“on-resonance”), and an integrated luminosity of 9.6 fb collected about 40 MeV below the $Y(4S)$ (“off-resonance”). In Ref. [10] we describe the silicon vertex tracker and drift chamber used for track and vertex reconstruction, the Cherenkov detector (DIRC), the electromagnetic calorimeter (EMC), and their use in particle identification (PID).

We reconstruct $B_{\rho h}^0$ candidates from combinations of two tracks and a π^0 candidate. We require that the PID of both tracks be inconsistent with the electron hypothesis, and the PID of the track used to form the ρ be inconsistent with the kaon hypothesis. The π^0 candidate mass must satisfy $0.11 < m(\gamma\gamma) < 0.16 \text{ GeV}/c^2$, where each photon is required to have an energy greater than 50 MeV in the laboratory frame and to exhibit an EMC cluster profile consistent with an electromagnetic shower. The mass of the ρ candidate must satisfy $0.4 < m(\pi^\pm \pi^0) < 1.3 \text{ GeV}/c^2$. To avoid most of the interference region, the B candidate is rejected if both the $\pi^+ \pi^0$ and $\pi^- \pi^0$ pairs satisfy this requirement. Taking advantage of the helicity structure of $B \rightarrow \rho h$ decays (h is denoted *bachelor track* hereafter), we require $|\cos \theta_\pi| > 0.25$, where θ_π is the angle between the π^0 momentum and the negative B momentum in the ρ rest frame. The bachelor track from the ρh decay must have a e^+e^- center-of-mass (c.m.) momentum above 2.4 GeV/ c .

For 86% of the $B^0 \rightarrow \rho h$ decays that pass the event selection, the pion from the ρ has momentum below this value, and thus the charge of the ρ is determined unambiguously. For the remaining events, the charge of the ρ is taken to be that of the selected $\pi^\pm \pi^0$ combination. With this procedure, 5% of the selected simulated signal events are assigned an incorrect charge.

To reject background from two-body B decays, the invariant masses of the $\pi^\pm h^\mp$ and $h^\pm \pi^0$ combinations must each be less than 5.14 GeV/ c^2 . Two kinematic variables are used to discriminate between signal- B decays and combinatorial background. One variable is the difference, ΔE , between the c.m. energy of the B candidate and $\sqrt{s}/2$, where \sqrt{s} is the total c.m. energy. The other variable is the beam-energy-substituted mass $m_{ES} \equiv \sqrt{(s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2/E_i^2 - \mathbf{p}_B^2}$, where the B momentum \mathbf{p}_B and the four-momentum of the initial state (E_i, \mathbf{p}_i) are defined in the laboratory frame. The ΔE distribution for $\rho\pi$ (ρK) signal peaks around 0 (-45) MeV since the

pion mass is always assigned to the bachelor track. We require $5.23 < m_{ES} < 5.29 \text{ GeV}/c^2$ and $-0.12 < \Delta E < 0.15 \text{ GeV}$, where the asymmetric ΔE window suppresses higher-multiplicity B background, which leads to mostly negative ΔE values. Discrimination between $\rho\pi$ and ρK events is provided by the Cherenkov angle θ_C and, to a lesser extent, by ΔE .

The dominant continuum background from $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) events is suppressed through the use of a neural network (NN) combining four discriminating variables: the reconstructed ρ mass, $\cos \theta_\pi$, and the two event-shape variables that are used in the Fisher discriminant of Ref. [2]. The NN is trained in the signal region with off-resonance data and simulated signal events. The final sample of signal candidates is selected with a cut on the NN output that retains $\sim 65\%$ (5%) of the signal (continuum).

Approximately 23% (20%) of simulated $\rho\pi$ (ρK) events have more than one ρh candidate passing the selection criteria. In these cases, we choose the candidate with the reconstructed π^0 mass closest to the nominal π^0 mass. A total of 20 497 events pass all selection criteria. The signal efficiency determined from Monte Carlo (MC) simulation is 20.7% (18.5%) for $\rho\pi$ (ρK) events; 31% (30%) of the selected events are misreconstructed, mostly due to combinatorial- π^0 background.

We use MC-simulated events to study the cross feed from other B decays. The charmless modes are grouped into 11 classes with similar kinematic and topological properties. Two additional classes account for the neutral and charged $b \rightarrow c$ decays. For each of the background classes, a component is introduced into the likelihood, with a fixed number of events. Backgrounds from two-, three-, and four-body decays to $\rho\pi$ are dominated by $B^+ \rightarrow \pi^+\pi^0$, $B^+ \rightarrow \rho^0\pi^+$, and longitudinally polarized $B^0 \rightarrow \rho^+\rho^-$ decays. The dominant two-body background for the ρK sample is $B^+ \rightarrow K^+\pi^0$, while for three- and four-body sources it is $B \rightarrow K^*\pi$ and higher kaon resonances, estimated from inclusive $B \rightarrow K\pi\pi$ measurements.

The time difference Δt is obtained from the measured distance between the z positions (along the beam direction) of the $B_{\rho h}^0$ and B_{tag}^0 decay vertices, and the boost $\beta\gamma = 0.56$ of the e^+e^- system [2,11]. To determine the flavor of the B_{tag}^0 we use the tagging algorithm of Ref. [11]. This produces four mutually exclusive tagging categories. We also retain untagged events in a fifth category to improve the efficiency of the signal selection and the sensitivity to charge asymmetries. Correlations between the B flavor tag and the charge of the reconstructed ρh candidate are observed in various B -background channels and evaluated with MC simulation.

We use an unbinned extended maximum likelihood fit to extract the $\rho\pi$ and ρK event yields, the CP parameters, and the other parameters defined in Eq. (1). The likelihood for the N_k candidates i tagged in category k is

$$\mathcal{L}_k = e^{-N'_k} \prod_{i=1}^{N'_k} \sum_h^{\pi, K} \left\{ N^{\rho h} \epsilon_k \mathcal{P}_{i,k}^{\rho h} + N_k^{q\bar{q},h} \mathcal{P}_{i,k}^{q\bar{q},h} + \sum_{j=1}^{N_B} \mathcal{L}_{ij,k}^{B,h} \right\}, \quad (2)$$

where N'_k is the sum of the signal and continuum yields (to be determined by the fit) and the fixed B -background yields, $N^{\rho h}$ is the number of signal events of type ρh in the entire sample, ϵ_k is the fraction of signal events tagged in category k , and $N_k^{q\bar{q},h}$ is the number of continuum background events with bachelor track of type h that are tagged in category k . The total likelihood \mathcal{L} is the product of likelihoods for each tagging category.

The probability density functions (PDFs) $\mathcal{P}_k^{\rho h}$, $\mathcal{P}_k^{q\bar{q},h}$ and the likelihood terms $\mathcal{L}_{j,k}^{B,h}$ are the product of the PDFs of five discriminating variables. The signal PDF is thus given by $\mathcal{P}_k^{\rho h} = \mathcal{P}_k^{\rho h}(m_{ES}) \cdot \mathcal{P}_k^{\rho h}(\Delta E) \cdot \mathcal{P}_k^{\rho h}(\text{NN}) \times \mathcal{P}_k^{\rho h}(\theta_C) \cdot \mathcal{P}_k^{\rho h}(\Delta t)$, where $\mathcal{P}_k^{\rho h}(\Delta t)$ contains the measured physics quantities defined in Eq. (1), diluted by the effects of mistagging and the Δt resolution. The PDF of the continuum contribution with bachelor track h is denoted $\mathcal{P}_k^{q\bar{q},h}$. The likelihood term $\mathcal{L}_{j,k}^{B,h} = N_{j,k}^{B,h} \mathcal{P}_{ij,k}^{B,h}$ corresponds to the B -background contribution j of the N_B classes with an expected event yield $N_{j,k}^{B,h}$.

The signal PDFs are decomposed into three parts with distinct distributions: signal events that are correctly reconstructed, misreconstructed signal events with right-sign ρ charge, and misreconstructed signal events with wrong-sign ρ charge. Their individual fractions are estimated by MC simulation. The m_{ES} , ΔE , and NN output PDFs for signal and B background are taken from the simulation except for the means of the signal Gaussian PDFs for m_{ES} and ΔE , which are free to vary in the fit. The continuum PDFs are described by six free parameters. The θ_C PDF is modeled as in Ref. [2]. The Δt -resolution function for signal and B -background events is a sum of three Gaussian distributions, with parameters determined from a fit to fully reconstructed B^0 decays [11]. The continuum Δt distribution is parametrized as the sum of three Gaussian distributions with common mean, two relative fractions, and three distinct widths that scale the Δt event-by-event error, yielding six free parameters. For continuum, two charge asymmetries and the ten parameters $N_k^{q\bar{q},h}$ are free. A total of 34 parameters, including signal yields and the parameters from Eq. (1), are varied in the fit.

The contributions to the systematic error on the signal parameters are summarized in Table I. The uncertainties associated with Δm_d and τ are estimated by varying these parameters within the uncertainty on the world average [12]. The uncertainties due to the signal model (PDF shapes, fraction of misreconstructed events) are obtained from a control sample of fully reconstructed $B^0 \rightarrow D^-\rho^+$ decays. We perform fits on large MC samples with the measured proportions of $\rho\pi/\rho K$ signal, and

TABLE I. Summary of the systematic uncertainties.

Error source	$N^{\rho K}$ (Events)	$N^{\rho\pi}$ (Events)	$A_{CP}^{\rho K}$	$A_{CP}^{\rho\pi}$	$C_{\rho\pi}$	$\Delta C_{\rho\pi}$	$S_{\rho\pi}$	$\Delta S_{\rho\pi}$
						(in units of 10^{-2})		
Δm_d and τ	0.1	0.1	0.0	0.0	0.4	0.4	0.2	0.1
Δt PDF	1.2	1.9	0.4	0.2	1.4	0.8	1.5	1.2
Signal model	4.0	13.1	1.2	0.8	0.7	0.8	1.4	1.0
Particle ID	0.6	0.7	0.5	0.2	0.1	0.1	0.1	0.1
Fit procedure	8.0	15.7	0.4	0.2	0.4	0.4	0.4	0.3
DCS decays	0.0	0.3	0.0	0.1	2.2	2.2	0.8	0.7
B background	16.0	14.2	7.9	2.8	3.0	3.5	2.1	1.8
Total	18.4	25.0	8.0	2.9	4.1	4.3	3.1	2.5

continuum and B backgrounds. Biases observed in these tests are due to imperfections in the PDF model; e.g., unaccounted correlations between discriminating variables. The biases are added in quadrature and assigned as a systematic uncertainty of the fit procedure. The systematic errors due to interference between the doubly-Cabibbo-suppressed (DCS) $\bar{b} \rightarrow \bar{u}c\bar{d}$ amplitude with the Cabibbo-favored $b \rightarrow c\bar{u}d$ amplitude for tagside B decays have been estimated from simulation by varying freely all relevant strong phases [13].

The main source of systematic uncertainty is the B -background model. The expected event yields from the background modes are varied according to the uncertainties in the measured or estimated branching fractions. Systematic errors due to possible nonresonant $B^0 \rightarrow \pi^+ \pi^- \pi^0$ decays are derived from experimental limits [5]. Repeating the fit without using the ρ -candidate mass and helicity angle gives results that are compatible with those reported here. Since B -background modes may exhibit CP violation, the corresponding parameters are varied within their physical ranges.

The maximum likelihood fit results in the event yields $N^{\rho\pi} = 428^{+34}_{-33}$ and $N^{\rho K} = 120^{+21}_{-20}$, where the errors are statistical. Correcting the yields by a small fit bias determined using the MC simulation (3% for $\rho\pi$ and 0% for




FIG. 1. Distributions of m_{ES} and ΔE for samples enhanced in $\rho\pi$ signal (a),(c) and ρK signal (b),(d). The solid curve represents a projection of the maximum likelihood fit result. The dashed curve represents the contribution from continuum events, and the dotted line indicates the combined contributions from continuum events and B -related backgrounds.

ρK), we find for the branching fractions

$$\begin{aligned} \mathcal{B}(B^0 \rightarrow \rho^\pm \pi^\mp) &= (22.6 \pm 1.8 \pm 2.2) \times 10^{-6}, \\ \mathcal{B}(B^0 \rightarrow \rho^- K^+) &= (7.3^{+1.3}_{-1.2} \pm 1.3) \times 10^{-6}, \end{aligned} \quad (3)$$

where the first errors are statistical and the second systematic. The systematic errors include an uncertainty of 7.7% for efficiency corrections, dominated by the uncertainty in the π^0 reconstruction efficiency. Figure 1 shows distributions of m_{ES} and ΔE , enhanced in signal content by cuts on the signal-to-continuum likelihood ratios of the other discriminating variables. Distributions of discriminating variables show satisfactory agreement with the fit. For the CP -violating parameters, we obtain

$$\begin{aligned} A_{CP}^{\rho\pi} &= -0.18 \pm 0.08 \pm 0.03, & A_{CP}^{\rho K} &= 0.28 \pm 0.17 \pm 0.08, \\ C_{\rho\pi} &= 0.36 \pm 0.18 \pm 0.04, & S_{\rho\pi} &= 0.19 \pm 0.24 \pm 0.03. \end{aligned} \quad (4)$$

For the other parameters in the description of the $B^0(\bar{B}^0) \rightarrow \rho\pi$ decay-time dependence, we find

$$\begin{aligned} \Delta C_{\rho\pi} &= 0.28^{+0.18}_{-0.19} \pm 0.04, \\ \Delta S_{\rho\pi} &= 0.15 \pm 0.25 \pm 0.03. \end{aligned}$$

We find the linear correlation coefficients $c_{C,\Delta C} = 0.18$ and $c_{S,\Delta S} = 0.23$, while all other correlations are smaller. As a validation of our treatment of the time dependence we allow τ and Δm_d to vary in the fit. We

find $\tau = (1.64 \pm 0.13) \text{ ps}$ and $\Delta m_d = (0.52 \pm 0.12) \text{ ps}^{-1}$; the remaining free parameters are consistent with the nominal fit. The raw time-dependent asymmetry $A_{B^0/\bar{B}^0} = (N_{B^0} - N_{\bar{B}^0})/(N_{B^0} + N_{\bar{B}^0})$ in the tagging categories dominated by kaons and leptons is represented in Fig. 2.

In summary, we have presented measurements of branching fractions and CP -violating asymmetries in $B^0 \rightarrow \rho^\pm \pi^\mp$ and $\rho^- K^+$ decays. We do not find evidence




FIG. 2. Time distributions for events selected to enhance the $\rho\pi$ signal tagged as (a) B_{tag}^0 and (b) \bar{B}_{tag}^0 , and (c) time-dependent asymmetry between B_{tag}^0 and \bar{B}_{tag}^0 . The solid curve is a likelihood projection of the fit result. The dashed line is the total B - and continuum-background contribution.

for either mixing-induced CP violation in the time-dependent asymmetry of $B^0 \rightarrow \rho^\pm \pi^\mp$ decays, or direct CP violation in $B^0 \rightarrow \rho^- K^+$. Our measurement of direct CP violation in $B^0 \rightarrow \rho^\pm \pi^\mp$ is consistent with zero within 2.0 standard deviations, when both statistical and systematic errors are taken into account.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway),

MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

*Also with Università di Perugia, Perugia, Italy.

†Also with Università della Basilicata, Potenza, Italy.

‡Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain.

§Deceased.

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 281802 (2002).
- [3] Belle Collaboration, K. Abe *et al.*, Phys. Rev. D **68**, 012001 (2003).
- [4] R. Aleksan, I. Dunietz, B. Kayser, and F. Le Diberder, Nucl. Phys. **B361**, 141 (1991).
- [5] *BABAR* Collaboration, B. Aubert *et al.*, BABAR-CONF-01-10, SLAC-PUB-8926, hep-ex/0107058 (2001).
- [6] Belle Collaboration, A. Gordon *et al.*, Phys. Lett. B **542**, 183 (2002); CLEO Collaboration, C. P. Jessop *et al.*, Phys. Rev. Lett. **85**, 2881 (2000).
- [7] H. J. Lipkin, Y. Nir, H. R. Quinn, and A. Snyder, Phys. Rev. D **44**, 1454 (1991).
- [8] A. E. Snyder and H. R. Quinn, Phys. Rev. D **48**, 2139 (1993).
- [9] *The BABAR Physics Book*, edited by P. F. Harrison and H. R. Quinn (Stanford Linear Accelerator Center, Stanford, CA, 1998).
- [10] *BABAR* Collaboration, A. Palano *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [11] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **66**, 032003 (2002); *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002).
- [12] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
- [13] O. Long, M. Baak, R. N. Cahn, and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).